



# An updated checklist of blowflies (Diptera: Calliphoridae) from India

Meenakshi Bharti

Department of Zoology, Punjabi University, Patiala, Punjab, India-147002.  
(email: adubharti@yahoo.co.in)

## Abstract

An updated checklist of blowflies (Diptera: Calliphoridae) is provided herewith. This has been carried due to some recent shufflings, new records, new species and erroneous placement of taxa in earlier reported lists. Now, the family Calliphoridae is represented by 9 subfamilies, 30 genera and 119 species from India.

**Keywords:** Blowflies, Calliphoridae, Diptera, India.

## Introduction

Since Senior-White's fauna of British India and Oriental Region (Diptera: Calliphoridae, Vol-VI) nothing much has been contributed on this family from India in terms of streamlining the taxonomy of this complicated group. Isolated inputs in the form of new species/records have appeared from national and international workers from time to time, but no data in term of number of species appeared since checklists by Nandi, 2004; Bharti, 2008 and Mitra and Sinha (published online). Unfortunately, the authors of online checklist have not consulted the literature pertaining to current status of various taxa, which have undergone shuffling in recent times thus leading to errors. Moreover, data needs to be updated due to recent shufflings/publications (Bharti and Kurahashi, 2009; Rognes, 2009 and Bharti and Kurahashi, 2010) and on the basis of recommendations by Rognes, 2010 (personal communication). For instance, as per Rognes, 2009 *Bengalia emarginatoides* is a new edition to India and Srilanka and *Bengalia pallidoxa* Seguy, 1946; *Bengalia chromatella* Seguy, 1946 and *Bengalia bezzi* Senior White, 1924 as junior synonyms of *Bengalia varicolor* (Fabricius, 1805). Rognes, 2010 (personal communication) also amended the status of species belonging to genus *Bengalia* which were erected by Lehrer, 2005. These include *Bengalia krishna* Lehrer, 2005 (=*B. martinleakei*), *Bengalia evanfoneae* Lehrer, 2005 (=*B. jejuna*), *Bengalia gandhana* Lehrer, 2005 (=*B. martinleakei*), *Bengalia indipyga* Lehrer, 2005 (=*B. kanoi*?).

In the wake of above mentioned discussion the updated list is as follows:

### Family: Calliphoridae

#### Subfamily: Ameniinae

##### Tribe: Catapicephalini

##### Genus: *Catapicephala* Macquart, 1850

*Catapicephala ingens* (Walker, 1860)

*Catapicephala pattoni* Senior-White, Aubertin & Smart, 1940

*Catapicephala micans* (Fabricius, 1805)

*Catapicephala splendens* Macquart, 1851

##### Tribe: Ameniini

##### Genus: *Silbomyia* Macquart, 1843

*Silbomyia asiatica* Crosskey, 1965

*Silbomyia parvula* Baranov, 1938

#### Subfamily: Helicoboscinae

##### Genus: *Gulmargia* Rognes, 1992

*Gulmargia angustisquama* Rognes, 1993

#### Subfamily: Calliphorinae

##### Tribe: Calliphorini

##### Genus: *Aldrichina* Townsend, 1934

*Aldrichina grahamni* (Aldrich, 1930)

##### Genus: *Calliphora* Robineau-Desvoidy, 1830

*Calliphora vicina* Robineau-Desvoidy, 1830

*Calliphora pattoni* Aubertin, 1931

*Callipora vomitoria* (Linneaus, 1758)

*Calliphora loewi* Enderlein, 1903

#### Tribe: Phumosiini

Genus: *Phumosia* Robineau-Desvoidy, 1830

*Phumosia indica* (Surcouf, 1914)

*Phumosia testacea* (Senior-White, 1923)

#### Subfamily: Bengaliinae

##### Tribe: Bengaliini

Genus: *Bengalia* Robineau-Desvoidy, 1830

*Bengalia emarginatoides* Rognes, 2009

*Bengalia labiata* Robineau-Desvoidy, 1830

*Bengalia hastativentris* Senior-White, 1923

*Bengalia jejuna* (Fabricius, 1787) (= *B. evanfoneae*, Lehrer, 2005)

*Bengalia jejuna* var. *quadrinotata* (Bigot, 1887)

*Bengalia lateralis* Macquart, 1843

*Bengalia martin-leakei* Senior-White, 1930 (= *B. Krishna* Lehrer, 2005, *B. gandhana* Lehrer, 2005)

*Bengalia escheri* Beazzi, 1913

*Bengalia xanthopyga* Senior-White, 1924

*Bengalia surcoufi* Senior-White, 1923

*Bengalia varicolor* (Fabricius, 1805) (= *B. beazzi* Senior-White, 1923, *B. pallidicoxa* Seguy, 1946, *B. chromatella* Seguy, 1946) Rognes, 2009

*Bengalia torosa* (Wiedemann, 1819)

#### Subfamily: Melanomyinae

Genus: *Melinda* Robineau-Desvoidy, 1830

*Melinda abdominalis* (Malloch, 1931)

*Melinda bengalensis* Nandi, 1994

*Melinda chambensis* Singh and Sidhu, 2007

*Melinda chandigarhensis* Singh and Sidhu, 2007

*Melinda pusilla* (Villeneuve, 1927) (= *Pollenia townsendi* Senior-White, Aubertin & Smart, 1940)

*Melinda pusilla indica* Kurahashi, 1970

*Melinda scutellata* (Senior-White, 1923)

#### Subfamily: Luciliinae

##### Tribe: Luciliini

Genus: *Hemipyrellia* Townsend, 1917

*Hemipyrellia ligurriens* (Wiedemann, 1830)

*Hemipyrellia pulchra* (Wiedemann, 1830)

Genus: *Hypopygiopsis* Townsend, 1916

*Hypopygiopsis infumata* (Bigot, 1877)

*Hypopygiopsis tumrasvini* Kurahashi, 1977

Genus: *Lucilia* Robineau-Desvoidy, 1830

*Lucilia ampullacea* Villeneuve, 1922

*Lucilia bazini* Seguy, 1934

*Lucilia calviceps* Beazzi, 1927 (New record, Bharti & Kurahashi, 2010)

*Lucilia cuprina* (Wiedemann, 1830)

*Lucilia illustris* (Meigen, 1826)

*Lucilia porphyrina* (Walker, 1856)

*Lucilia papuensis* Macquart, 1843

*Lucilia sericata* (Meigen, 1826)

#### Subfamily: Polleniinae

##### Tribe: Polleniini

Genus: *Pollenia* Robineau-Desvoidy, 1830

*Pollenia hazarae* (Senior-White, 1923)

*Pollenia rudis* (Fabricius, 1794)

##### Tribe: Melanodexiini

Genus: *Morinia* Robineau-Desvoidy, 1830

*Morinia argenticincta* (Senior-White, 1923)

Genus: *Wilhelmina* Schmidt & Villeneuve, 1932

*Wilhelmina indica* Sidhu & Singh, 2005

Genus: *Polleniopsis* Townsend, 1917

*Polleniopsis asiatica* (Senior-White, 1923) new comb. (Kurahashi, 1972)

*Polleniopsis kashmirensis* Kurahashi & Okadome, 1976

*Polleniopsis pilosa* (Townsend, 1917)

Genus: *Tainanina* Villeneuve, 1926

*Tainanina pilisquama* (Senior-White, 1925)

*Tainanina sarcophagooides* (Malloch, 1931)

Genus: *Onesia* Robineau-Desvoidy, 1830

*Onesia khasiensis* (Senior-White, 1922)

Genus: *Dexopollenia* Townsend, 1917

*Dexopollenia flava* Aldrich, 1930

*Dexopollenia testacea* (Townsend, 1917)

#### Subfamily: Chrysomyinae

##### Tribe: Chrysomyini

Genus: *Chrysomya* Robineau-Desvoidy, 1830

*Chrysomya albiceps* (Wiedemann, 1819)

*Chrysomya bezziana* Villeneuve, 1914

*Chrysomya defixa* (Walker, 1857)

*Chrysomya indica* Sinha and Nandi, 2004

*Chrysomya megacephala* (Fabricius, 1794) (fdf/sdf, two forms in India, Bharti & Kurahashi, 2009)

*Chrysomya nigripes* Aubertin, 1932

*Chrysomya pinguis* (Walker, 1858)

*Chrysomya phaonis* (Seguy, 1928)

*Chrysomya rufifacies* (Macquart, 1843)

*Chrysomya villeneuvi* Patton, 1922

#### Subfamily: Rhiniinae

##### Tribe: Rhiniini

###### Genus: *Chlororhinia* Townsend, 1917

*Chlororhinia exempta* (Walker, 1856)

*Chlororhinia tamenensis* Joseph & Rao, 1972

###### Genus: *Idiella* Braeuer & Berensteamn, 1889

*Idiella divisa* (Walker, 1861)

*Idiella euidielloides* Senior-White, 1923

*Idiella mandarina* (Wiedemann, 1830)

*Idiella tripartia* (Bigot, 1874)

###### Genus: *Idiellopsis* Townsend, 1917

*Idiellopsis xanthogaster* (Wiedemann, 1820)

###### Genus: *Rhinia* Robineau-Desvoidy, 1830

*Rhinia mallochi* (Senior-White, Aubertin & Smart, 1940)

*Rhinia apicalis* (Wiedemann, 1830)

###### Genus: *Stomorrhina* Rondani, 1861

*Stomorrhina discolor* (Fabricius, 1794)

*Stomorrhina lunata* (Fabricius, 1805)

*Stomorrhina melastoma* (Wiedemann, 1830)

*Stomorrhina procula* (Walker, 1849)

*Stomorrhina simplex* (Walker, 1857)

*Stomorrhina townsendi* Kurahashi, 1997

*Stomorrhina unicolor* (Macquart, 1851)

*Stomorrhina xanthogaster* (Wiedemann, 1820)

##### Tribe: Cosminini

###### Genus: *Cosmina* Robineau-Desvoidy, 1830

*Cosmina bicolor* (Walker, 1856)

*Cosmina limbipennis* (Macquart, 1848)

*Cosmina prasina* (Brauer & Bergenstamm, 1889)

*Cosmina viridis* Townsend, 1917

*Cosmina simplex* (Walker, 1858)

###### Genus: *Thoracites* Brauer & Bergenstamm, 1891

*Thoracites abdominalis* (Fabricius, 1805)

##### Tribe: Isomyiini

###### Genus: *Isomyia* Walker, 1860

*Isomyia aurifacies* James, 1970

*Isomyia delectans* Walker, 1860

*Isomyia electa* Villeneuve, 1927

*Isomyia fulvicornis* (Bigot, 1887)

*Isomyia fuscocincta* Bigot, 1877

*Isomyia gomezmenori* (Peris, 1951)

*Isomyia nebulosa* (Townsend, 1917)

*Isomyia nigrofasciata* (Peris, 1951)

*Isomyia oestracea* (Seguy, 1934)

*Isomyia perisi* James, 1970

*Isomyia phryxe* Seguy, 1949

*Isomyia pseudonepalana* (Senior-White, Aubertin & Smart, 1940)

*Isomyia pseudoviridana* (Peris, 1952)

*Isomyia sivah* (Bigot, 1878)

*Isomyia versicolor* (Bigot, 1877)

*Isomyia viridaurea* (Wiedemann, 1819)

*Isomyia zeylanica* Senior-White, Aubertin & Smart, 1940

###### Genus: *Strongyloneura* Bigot, 1886

*Strongyloneura nudus* Ghezta and Kumar, 1991

*Strongyloneura prolata* (Walker, 1860)

##### Tribe: Rhyncomyiini

###### Genus: *Borbororhinia* Townsend, 1917

*Borbororhinia bivittata* (Walker, 1856)

###### Genus: *Metallea* van der Wulp, 1880

*Metallea notata* van der Wulp, 1880

*Metallea flavibasis* Senior-White, 1922

*Metallea setosa* (Townsend, 1917)

###### Genus: *Rhyncomya* Robineau-Desvoidy, 1830

*Rhyncomya callopis* Loew, 1856

*Rhyncomya catalypsa* Seguy, 1946

*Rhyncomya divisa* (Walker, 1856)

*Rhyncomya flavipes* Robineau-Desvoidy, 1830

*Rhyncomya pollinosa* (Townsend, 1917)

*Rhyncomya sarotes* Seguy, 1928

*Rhyncomya viridaurea* Wiedemann, 1824

##### Doubtful Species

*Bengalia rivanella* Lehrer, 2005

*Bengalia shivanella* Lehrer, 2005

To sum up, family Calliphoridae is represented by 120 species from India, out of which 22 species are endemic to this region.

## Acknowledgements

Financial assistance rendered by Department of Science and Technology, New Delhi vide letter number SR/WOS-A/LS-20/2009 is greatly acknowledged.

## References

- Bharti, M. 2008. Diptera: Calliphoridae, Rhiniidae from India. *Journal of Entomological Research* 32 (1): 79-82.
- Bharti, M. and Kurahashi, H. 2009. Finding of feral derived form (fdf) of *Chrysomya megacephala* (Fabricius) from India with an evolutionary novelty (Diptera: Calliphoridae). *Japanese Journal of Systematic Entomology* 15 (2): 411-413.
- Bharti, M. and Kurahashi, H. 2010. *Lucilia calviceps* Bezzii, new record from India (Diptera: Calliphoridae). *Halteres* 2: 29-30.
- Kurahashi, H. 1970. Tribe Calliphorini from Australian and Oriental regions, 1. *Melinda*-group (Diptera: Calliphoridae). *Pacific Insects* 12 (3): 519-542.
- Kurahashi, H. 1972. Tribe Calliphorini from Australian and Oriental region, iv. *Onesia*-group: genus *Polleniopsis* (Diptera: Calliphoridae). *Pacific insects* 14 (4):709-724.
- Kurahashi, H. 1977. The tribe Luciliini from Australian and Oriental regions 1. Genus *Hypopygiopsis* Townsend (Diptera: Calliphoridae). *Kontyu* 45(4): 553-562.
- Nandi, B. C. 2004. Checklist of Calliphoridae (Diptera) of India. *Records of the Zoological Survey of India, Occasional paper No. 231:* 1-47.
- Rognes, K. 2009. Revision of the oriental species of *Bengalia peuhui* species group (Diptera, Calliphoridae). *Zootaxa* 2251: 1-76.
- Senior- White, R., Aubertin,D. and Smart, J. 1940. The Fauna of British India, including the remainder of the oriental region, Diptera Vol.VI. Family Calliphoridae. London: Taylor and Francis.
- Sidhu, I.S. and Singh, D. 2005. First record of the genus *Wilhelmina* Schmitz and Villeneuve from India with description of a new species (Diptera: Calliphoridae). *Entomon* 30 (3): 255-259.
- Singh, D. and Sidhu, I.S. 2004a. New records of blowflies (Diptera: Calliphoridae) from India. *Entomon* 29 (2): 203-206.
- Singh, D. and Sidhu, I.S. 2004b. A checklist of blowflies (Diptera: Calliphoridae) from North-West of India. *Uttar Pradesh Journal of Zoology* 24 (1): 63-71.
- Singh, D. and Sidhu, I.S. 2007. Two new species of *Melinda* Robineau-Desvoidy (Diptera: Calliphoridae) from India with a key to Indian species of this genus. *Journal of Bombay Natural History Society* 104 (1): 55-57.